

Regularity conditions for arbitrary Leavitt path algebras

Gene Abrams · Kulumani M. Rangaswamy

Received: date / Accepted: date

Abstract We show that if E is an arbitrary acyclic graph then the Leavitt path algebra $L_K(E)$ is locally K -matricial; that is, $L_K(E)$ is the direct union of subalgebras, each isomorphic to a finite direct sum of finite matrix rings over the field K . (Here an *arbitrary* graph means that neither cardinality conditions nor graph-theoretic conditions (e.g. row-finiteness) are imposed on E . These unrestrictive conditions are in contrast to the hypotheses used in much of the literature on this subject.) As a consequence we get our main result, in which we show that the following conditions are equivalent for an arbitrary graph E : (1) $L_K(E)$ is von Neumann regular. (2) $L_K(E)$ is π -regular. (3) E is acyclic. (4) $L_K(E)$ is locally K -matricial. (5) $L_K(E)$ is strongly π -regular. We conclude by showing how additional regularity conditions (unit regularity, strongly clean) can be appended to this list of equivalent conditions.

Keywords Leavitt path algebra; acyclic graph; von Neumann regular algebra

Mathematics Subject Classification (2000) Primary: 16S99, 16E50. Secondary: 16W50, 46L89

In this article we investigate Leavitt path algebras $L_K(E)$ over an arbitrary directed graph E and an arbitrary field K . Here we require no restriction on either the size of the graph (i.e., vertex and edge sets may be of any cardinality), or on graph-theoretic constraints (i.e., no row-finiteness conditions are assumed). Our goal is to classify in ring-theoretic terms the Leavitt path algebras of the form $L_K(E)$ where E is acyclic. A useful tool in our study is a construction presented in Proposition 1, which enables us to realize $L_K(E)$ as a directed union of subalgebras, each of which is a Leavitt path algebra of a suitable finite graph. An effective use of Proposition 1 leads to our main result, Theorem 1, in which we establish the equivalence of these conditions for an arbitrary directed graph E and field K : (1) $L_K(E)$ is von Neumann regular. (2)

Gene Abrams and Kulumani M. Rangaswamy
Department of Mathematics
University of Colorado at Colorado Springs
Colorado Springs, Colorado, 80933 U.S.A.
E-mail: abrams@math.uccs.edu krangasw@uccs.edu
Corresponding author: G. Abrams 7192623182(v) 7192623605(f)

$L_K(E)$ is π -regular. (3) E is acyclic. (4) $L_K(E)$ is locally K -matricial (i.e., $L_K(E)$ is a direct union of subalgebras, each of which is isomorphic to a finite direct sum of finite matrix rings over K). (5) $L_K(E)$ is strongly π -regular. We conclude the article by discussing various additional ring-theoretic conditions which in the context of Leavitt path algebras are equivalent to E being acyclic.


We begin by giving a terse reminder of the germane definitions. For a more complete description and discussion, see e.g. [2] or [8]. A (directed) graph $E = (E^0, E^1, r, s)$ consists of two sets E^0, E^1 and functions $r, s : E^1 \rightarrow E^0$. (The sets E^0 and E^1 are allowed to be of arbitrary cardinality.) The elements of E^0 are called *vertices* and the elements of E^1 *edges*. A *path* μ in a graph E is a sequence of edges $\mu = e_1 \dots e_n$ such that $r(e_i) = s(e_{i+1})$ for $i = 1, \dots, n-1$. In this case, $s(\mu) := s(e_1)$ is the *source* of μ , $r(\mu) := r(e_n)$ is the *range* of μ , and n is the *length* of μ . We view the elements of E^0 as paths of length 0. If $\mu = e_1 \dots e_n$ is a path in E , and if $v = s(\mu) = r(\mu)$ and $s(e_i) \neq s(e_j)$ for every $i \neq j$, then μ is called a *cycle based at v*. If $s^{-1}(v)$ is a finite set for every $v \in E^0$, then the graph E is called *row-finite*.

Definition 1 Let E be any directed graph, and K any field. The *Leavitt path K -algebra* $L_K(E)$ of E with coefficients in K is the K -algebra generated by a set $\{v \mid v \in E^0\}$ of pairwise orthogonal idempotents, together with a set of variables $\{e, e^* \mid e \in E^1\}$, which satisfy the following relations:

- (1) $s(e)e = er(e) = e$ for all $e \in E^1$.
- (2) $r(e)e^* = e^*s(e) = e^*$ for all $e \in E^1$.
- (3) (CK1) $e^*e' = \delta_{e,e'}r(e)$ for all $e, e' \in E^1$.
- (4) (CK2) $v = \sum_{\{e \in E^1 \mid s(e)=v\}} ee^*$ for every vertex $v \in E^0$ having $1 \leq |s^{-1}(v)| < \infty$.

For any $F \subseteq E^1$ the set $\{e^* \mid e \in F\}$ will be denoted by F^* . We let $r(e^*)$ denote $s(e)$, and we let $s(e^*)$ denote $r(e)$. If $\mu = e_1 \dots e_n$ is a path, then we denote by μ^* the element $e_n^* \dots e_1^*$ of $L_K(E)$.

Many well-known algebras arise as the Leavitt path algebra of a graph. For instance, the classical Leavitt algebras $L_K(1, n)$ for $n \geq 2$ arise as the algebras $L_K(R_n)$ where R_n is the “rose with n petals” graph described in Example 1 below. (See e.g. [1, Section 3].) Also, for each $n \in \mathbb{N} = \{1, 2, \dots\}$, the full matrix ring $M_n(K)$ arises as the Leavitt path algebra of the oriented n -line graph



while the Laurent polynomial ring $K[x, x^{-1}]$ arises as the Leavitt path algebra of the “one vertex, one loop” graph



A (possibly nonunital) ring R is called a *ring with local units* in case for each finite subset $S \subseteq R$ there is an idempotent $e \in R$ with $S \subseteq eRe$. If E is a graph for which E^0 is finite then we have $\sum_{v \in E^0} v$ is the multiplicative identity in $L_K(E)$; otherwise, $L_K(E)$ is a ring with a set of local units consisting of sums of distinct vertices. Conversely, if $L_K(E)$ is unital, then E^0 is finite. $L_K(E)$ is a $\mathbb{Z}$ -graded K -algebra, spanned as a K -vector space by $\{pq^* \mid p, q \text{ are paths in } E\}$. (Recall that the elements of E^0 are viewed as paths of length 0, so that this set includes elements of the form v with $v \in E^0$.) In

particular, for each $n \in \mathbb{Z}$, the degree n component $L_K(E)_n$ is spanned by elements of the form $\{pq^* \mid \text{length}(p) - \text{length}(q) = n\}$. The degree of an element x , denoted $\deg(x)$, is the lowest number n for which $x \in \bigoplus_{m \leq n} L_K(E)_m$. The K -linear extension of the assignment $pq^* \mapsto qp^*$ (for p, q paths in E) yields an involution on $L_K(E)$, which we denote simply as $*$.

A subgraph G of a graph E is called *complete* in case, for each $v \in G^0$ having $1 \leq |s_G^{-1}(v)| < \infty$, we have $s_G^{-1}(v) = s_E^{-1}(v)$. (In other words, a subgraph G of E is complete if, whenever $v \in G^0$ emits a nonzero, finite number of edges in G , then necessarily the subgraph G contains all of the edges in E emitted by v .) The natural inclusion map $L_K(G) \hookrightarrow L_K(E)$ is a ring homomorphism precisely when G is a complete subgraph of E , so that complete subgraphs of E naturally give rise to subalgebras of $L_K(E)$. One of our main objectives in this article is to show how to construct subalgebras of $L_K(E)$ which need not arise in this way. This in turn will allow us to describe algebras of the form $L_K(E)$ as unions of subalgebras possessing various ring-theoretic properties, even in situations where E lacks complete subgraphs possessing corresponding graph-theoretic properties. We achieve this objective in Proposition 1. The construction is based on an idea presented by Raeburn and Szymański in [13, Definition 1.1]; this work was brought to our attention by E. Pardo.

Definition 2 Let E be a graph, and let F be a finite set of edges in E . We define $s(F)$ (resp. $r(F)$) to be the sets of those vertices in E which appear as the source (resp. range) vertex of at least one element of F . We define a graph E_F as follows:

$$E_F^0 = F \cup (r(F) \cap s(F) \cap s(E^1 \setminus F)) \cup (r(F) \setminus s(F)),$$

$$E_F^1 = \{(e, f) \in F \times E_F^0 \mid r(e) = s(f)\} \cup \{(e, r(e)) \mid e \in F \text{ with } r(e) \in (r(F) \setminus s(F))\},$$


$$\text{and where } s((x, y)) = x, \quad r((x, y)) = y \quad \text{for any } (x, y) \in E_F^1.$$

Note that, since F is finite, the graph E_F is finite (regardless of the size of E).

Remarks: 1. It is conventional to define $s(v) = v$ for each vertex v in E . Because of that, the expression in rectangular brackets in Definition 2 for E_F^1 is redundant. However, we choose to keep this expression in the definition, as it makes the correspondence between E_F^1 and the set G^1 in the proof of Proposition 1 more transparent.

2. While the construction presented in Definition 2 is similar to that given in [13, Definition 1.1], there are indeed some significant differences. For instance, the construction of [13, Definition 1.1] requires that the graph E has no sinks, while the construction presented here has no such stipulation. Additionally, even in situations where E is a graph with no sinks and F is a finite subset of E^1 , the two constructions can in fact yield different corresponding graphs E_F . However, the underlying goal of each of the two constructions is the same, namely, to produce a subalgebra of a graph algebra which is isomorphic to the graph algebra of a finite graph.

Example 1 For clarity, we provide an example of the graph E_F constructed in the previous definition. Let E be the “rose with n -petals” graph




Let $F = \{y_1\}$. Then $E_F^0 = \{y_1\} \cup \{v\}$, and $E_F^1 = \{(y_1, y_1), (y_1, v)\}$. Pictorially, E_F is given by

$$E_F = (y_1, y_1) \curvearrowright \bullet_{y_1} \xrightarrow{(y_1, v)} \bullet_v$$

This example indicates that various properties of the graph E need not pass to the graph E_F . For instance, E is cofinal, while E_F is not. In particular, $L_K(E)$ is a simple algebra, while $L_K(E_F)$ is not. (See [2] for a more complete discussion.)

Our interest in the construction given in Definition 2 can be generally described as follows. We seek to place each finite set of elements taken from the Leavitt path algebra $L_K(E)$ inside a subalgebra of $L_K(E)$ which possesses certain 'finiteness' properties. In case E is row-finite, by [5, Lemma 3.2] we can realize $L_K(E)$ as the direct union of subalgebras of the form $L_K(E_i)$ where each E_i is a finite, complete subgraph of E . In the general case, however, we need not have such a description of $L_K(E)$. For instance, if $\aleph$ is an infinite cardinal, and $\text{Clock}(\aleph)$ denotes the 'infinite clock' graph



having $\aleph$ edges, then there are *no* nontrivial finite complete subgraphs of $\text{Clock}(\aleph)$.

Example 2 It will be instructive to consider the E_F construction of Definition 2 within the infinite clock graph $E = \text{Clock}(\aleph)$. So let v denote the center vertex, let f denote one of the edges, and let w denote $r(f)$. Let $F = \{f\}$. Then $E_F^0 = \{f\} \cup \{w\}$, while $E_F^1 = \{(f, w)\}$. Thus E_F is the graph

$$E_F = \bullet_f \xrightarrow{(f, w)} \bullet_w$$

with two vertices, and one edge connecting them. In particular, $L(E_F) \cong M_2(K)$.

Although in general E_F need not be a subgraph of E (indeed, as seen in Example 1, E_F may contain more vertices than does E), there is an important relationship between the Leavitt path algebras $L_K(E_F)$ and $L_K(E)$, as we now show.

Proposition 1 *Let F be a finite set of edges in a graph E . Then there is an algebra homomorphism $\theta : L_K(E_F) \rightarrow L_K(E)$ having the properties:*

- (1) $F \cup F^* \subseteq \text{Im}(\theta)$.
- (2) If $w \in r(F)$, then $w \in \text{Im}(\theta)$.
- (3) If $w \in E^0$ has $s_E^{-1}(w) \neq \emptyset$ and $s_E^{-1}(w) \subseteq F$, then $w \in \text{Im}(\theta)$.

Proof We define subsets G^0 and G^1 of $L_K(E)$ as follows.

$$\begin{aligned} G^0 &= \{ee^* \mid e \in F\} \\ &\cup \{v - \sum_{f \in F, s(f)=v} ff^* \mid v \in r(F) \cap s(F) \cap s(E^1 \setminus F)\} \\ &\cup \{v \mid v \in r(F) \setminus s(F)\} \end{aligned}$$

and

$$\begin{aligned} G^1 = & \{eff^* \mid e, f \in F, s(f) = r(e)\} \\ & \cup \{e - \sum_{f \in F, s(f)=r(e)} eff^* \mid r(e) \in r(F) \cap s(F) \cap s(E^1 \setminus F)\} \\ & \cup \{e \in F \mid r(e) \in r(F) \setminus s(F)\}. \end{aligned}$$

We define $\theta : L_K(E_F) \rightarrow L_K(E)$ as follows.

There are three different types of vertices in E_F . If $w \in E_F^0$ has form $w = e \in F$, then define

$$\theta(w) = ee^*.$$

If $w \in E_F^0$ has form $w = v$ with $v \in r(F) \cap s(F) \cap s(E^1 \setminus F)$, then define

$$\theta(w) = v - \sum_{f \in F, s(f)=v} ff^*.$$

If $w \in E_F^0$ has form $w = v$ with $v \in r(F) \setminus s(F)$, then define

$$\theta(w) = w.$$

Note that in each case we have $\theta(w) \in G^0$.

There are three different types of edges in E_F . If $h \in E_F^1$ has form $h = (e, f)$ with $f \in F$, then define

$$\theta(h) = eff^*.$$

If $h \in E_F^1$ has form $h = (e, r(e))$ with $r(e) \in r(F) \cap s(F) \cap s(E^1 \setminus F)$, then define

$$\theta(h) = e - \sum_{f \in F, s(f)=r(e)} eff^*.$$

If $h \in E_F^1$ has form $h = (e, r(e))$ with $r(e) \in r(F) \setminus s(F)$, then define

$$\theta(h) = e.$$

Note that in each case we have $\theta(h) \in G^1$.

For each $h \in E_F^1$ we define $\theta(h^*) = (\theta(h))^*$ in $L_K(E)$.

It is now a long, straightforward check to verify that θ is compatible with the four types of relations which define $L_K(E_F)$ (refer to Definition 1). As a representative example of the computations required here, we offer the following. Let $w \in E_F^0$ have the form $w = e \in F$. Then the (CK2) relation at e in $L_K(E_F)$ is the equation

$$\sum_{g \in E_F^1, s(g)=e} gg^* = e.$$

But $s(g) = e$ in E_F^1 means $g = (e, f)$ where either $f \in F$ has $s(f) = r(e)$, or $g = (e, r(e))$ with $r(e) \in r(F) \cap s(F) \cap s(E^1 \setminus F)$, or $g = (e, r(e))$ with $r(e) \in r(F) \setminus s(F)$. So the (CK2) relation at e in $L_K(E_F)$ takes the form

$$\begin{aligned}
e = & \sum_{f \in F, s(f)=r(e)} (e, f)(e, f)^* + \sum_{w \in r(F) \cap s(F) \cap s(E^1 \setminus F), w=r(e)} (e, w)(e, w)^* \\
& + \sum_{w \in r(F) \setminus s(F), w=r(e)} (e, w)(e, w)^*.
\end{aligned}$$

Note that empty sums are interpreted as 0. Also, the final two summation expressions are in fact either singletons or empty, depending on whether $r(e) \in r(F) \cap s(F) \cap s(E^1 \setminus F)$ or $r(e) \in r(F) \setminus s(F)$.

We must show that the corresponding equation under θ holds in $L_K(E)$. In other words, we must show

$$\begin{aligned}
ee^* = & \sum_{f \in F, s(f)=r(e)} (eff^*)(eff^*)^* \\
& + \sum_{w \in r(F) \cap s(F) \cap s(E^1 \setminus F), w=r(e)} [e - \sum_{f \in F, s(f)=r(e)} eff^*][e - \sum_{f \in F, s(f)=r(e)} eff^*]^* \\
& + \sum_{w \in r(F) \setminus s(F), w=r(e)} ee^*.
\end{aligned}$$

There are two cases. If $r(e) \in r(F) \setminus s(F)$, then this equation simply becomes $ee^* = ee^*$ and we are done. On the other hand, if $r(e) \in s(F)$, then note the second 'sum' $\sum_{w \in r(F) \cap s(F) \cap s(E^1 \setminus F), w=r(e)} [e - \sum_{f \in F, s(f)=r(e)} eff^*][e - \sum_{f \in F, s(f)=r(e)} eff^*]^*$ is in fact simply the single expression $[e - \sum_{f \in F, s(f)=r(e)} eff^*][e - \sum_{f \in F, s(f)=r(e)} eff^*]^*$. So the right hand side is

$$\begin{aligned}
& \sum_{f \in F, s(f)=r(e)} (eff^*)ff^*e^* + [e - \sum_{f \in F, s(f)=r(e)} eff^*][e - \sum_{f \in F, s(f)=r(e)} eff^*]^* \\
= & \sum_{f \in F, s(f)=r(e)} eff^*e^* + [ee^* - \sum_{f \in F, s(f)=r(e)} eff^*e^*] \quad (\text{by (CK1) and (CK2)}) \\
= & ee^*.
\end{aligned}$$

In a similar manner one can verify the compatibility of θ with all the remaining relations which define $L_K(E_F)$. Thus we conclude that θ extends to an algebra homomorphism

$$\theta : L_K(E_F) \rightarrow L_K(E).$$

By definition we have $\text{Im}(\theta)$ is the subalgebra of $L_K(E)$ generated by $G^0, G^1, (G^1)^*$. It will be helpful later to note that for each $x \in G^1 \cup (G^1)^*$ there exist $y, y' \in G^0$ for which $xyy' = x$. In particular, if an element $z \in L_K(E)$ is orthogonal to every element of G^0 , then necessarily z is orthogonal to every element in $\text{Im}(\theta)$.

We are now in position to verify the three claimed properties of $\text{Im}(\theta)$. For (1), we show that every $f \in F$ is contained in $\text{Im}(\theta)$. Suppose first that $f \in F$ has $s_E^{-1}(r(f)) \subseteq$

F . If $s_E^{-1}(r(f)) = \emptyset$, then $r(f) \in r(F) \setminus s(F)$ vacuously, so by definition $f \in G^1 \subseteq \text{Im}(\theta)$. On the other hand, if $s_E^{-1}(r(f)) \neq \emptyset$, then we have $fgg^* \in G^1$ for all $g \in F$ having $r(f) = s(g)$. But then by hypothesis this is the same as the collection of $g \in E^1$ having $r(f) = s(g)$. Thus we have $\{fgg^* \mid g \in E^1, s(g) = r(f)\} \subseteq \text{Im}(\theta)$, so that in particular $\text{Im}(\theta)$ contains $\sum_{g \in E^1, s(g)=r(f)} fgg^* = f \cdot \sum_{g \in E^1, s(g)=r(f)} gg^* = f \cdot r(f) = f$.

On the other hand, suppose that $f \in F$ has the property that $s_E^{-1}(r(f)) \not\subseteq F$. Then there are two possibilities. In the first case, $s_E^{-1}(r(f)) \neq \emptyset$ and $s_E^{-1}(r(f)) \cap F = \emptyset$. (In other words, there are edges in E which are emitted from $r(f)$, but none of these edges are in F .) But then $r(f) \notin s(F)$, so that $f \in G^1$ by definition, so that $f \in \text{Im}(\theta)$. In the second case, suppose $s_E^{-1}(r(f)) \cap F \neq \emptyset$. Then either we have $s_E^{-1}(r(f)) \subseteq F$ (in which case we are done by the previous paragraph), or we have $r(f) \in r(F) \cap s(F) \cap s(E^1 \setminus F)$. In this situation we have $fgg^* \in G^1 \subseteq \text{Im}(\theta)$ for all $g \in F$ having $s(g) = r(f)$, so we in particular have $\sum_{g \in F, s(g)=r(f)} fgg^*$ in $\text{Im}(\theta)$ as well. But by definition we also have the element $f - \sum_{g \in F, s(g)=r(f)} fgg^*$ in $G^1 \subseteq \text{Im}(\theta)$. Then

$$f = \left(\sum_{g \in F, s(g)=r(f)} fgg^* \right) + \left(f - \sum_{g \in F, s(g)=r(f)} fgg^* \right) \in \text{Im}(\theta).$$

Thus we conclude that $F \subseteq \text{Im}(\theta)$. But for each $x \in \text{Im}(\theta)$ we have $x^* \in \text{Im}(\theta)$ by definition. Thus $F \cup F^* \subseteq \text{Im}(\theta)$, thereby establishing (1).

In particular, if $w = r(f)$ for $f \in F$, then $w = f^*f \in \text{Im}(\theta)$, yielding (2). For (3), suppose $s^{-1}(w) \neq \emptyset$, and $s^{-1}(w) \subseteq F$. Then each ff^* for $f \in E^1$ having $s(f) = w$ is in G^0 , so that $\sum_{f \in E^1, s(f)=w} ff^*$ is in $\text{Im}(\theta)$. But this last sum is precisely w by (CK2).

We remark here that for θ as given in Proposition 1, $\theta(w) \neq 0$ for all three possible types of $w \in E_F^0$. (That $\theta(w) \neq 0$ in case $w \in r(F) \cap s(F) \cap s(E^1 \setminus F)$ hinges on the fact that there exists $g \in E^1 \setminus F$ having $s(g) = w$.) This in turn will allow us to conclude, in certain situations (including the situation where E is acyclic), that θ is in fact a monomorphism. (See e.g. [1].) However, we will not utilize this additional property of θ in the sequel.

With Proposition 1 in hand, we now construct the subalgebras of $L_K(E)$ which play the central role in our main result, Theorem 1.

The Subalgebra Construction

Let E be any graph, K any field, and $\{a_1, a_2, \dots, a_\ell\}$ any finite subset of nonzero elements of $L_K(E)$. For each $1 \leq r \leq \ell$ write

$$a_r = k_{c_1} v_{c_1} + k_{c_2} v_{c_2} + \dots + k_{c_{j(r)}} v_{c_{j(r)}} + \sum_{i=1}^{t(r)} k_{r_i} p_{r_i} q_{r_i}^*$$

where each k_j is a nonzero element of K , and, for each $1 \leq i \leq t(r)$, at least one of p_{r_i} or q_{r_i} has length at least 1. (That such a representation for each a_r exists follows from properties of $L_K(E)$ mentioned previously.) Let F denote the (necessarily finite) set of

those edges in E which appear in the representation of some p_{r_i} or q_{r_i} , $1 \leq r_i \leq t(r)$, $1 \leq r \leq \ell$. Now consider the set

$$S = \{v_{c_1}, v_{c_2}, \dots, v_{c_{j(r)}} \mid 1 \leq r \leq \ell\}$$

of vertices which appear in the displayed description of a_r for some $1 \leq r \leq \ell$. We partition S into subsets as follows:

$$S_1 = S \cap r(F),$$

and, for the remaining vertices $T = S \setminus S_1$, we define

$$S_2 = \{v \in T \mid s_E^{-1}(v) \subseteq F \text{ and } s_E^{-1}(v) \neq \emptyset\}$$

$$S_3 = \{v \in T \mid s_E^{-1}(v) \cap F = \emptyset\}$$

$$S_4 = \{v \in T \mid s_E^{-1}(v) \cap F \neq \emptyset \text{ and } s_E^{-1}(v) \cap (E^1 \setminus F) \neq \emptyset\}.$$

Let E_F be the graph as constructed in Definition 2 corresponding to this set F , and let $\theta : L_K(E_F) \rightarrow L_K(E)$ be the homomorphism described in Proposition 1.

Definition 3 Let E be any graph, K any field, and $\{a_1, a_2, \dots, a_\ell\}$ any finite subset of nonzero elements of $L_K(E)$. Consider the notation presented in The Subalgebra Construction. We define $B(a_1, a_2, \dots, a_\ell)$ to be the K -subalgebra of $L_K(E)$ generated by the set $\text{Im}(\theta) \cup S_3 \cup S_4$. That is,

$$B(a_1, a_2, \dots, a_\ell) = \langle \text{Im}(\theta), S_3, S_4 \rangle.$$

Proposition 2 Let E be any graph, K any field, and $\{a_1, a_2, \dots, a_\ell\}$ any finite subset of nonzero elements of $L_K(E)$. Let F denote the subset of E^1 presented in The Subalgebra Construction. For $w \in S_4$ let u_w denote the element $w - \sum_{f \in F, s(f)=w} f f^*$ of $L_K(E)$. Then

- (1) $\{a_1, a_2, \dots, a_\ell\} \subseteq B(a_1, a_2, \dots, a_\ell)$.
- (2) $B(a_1, a_2, \dots, a_\ell) = \text{Im}(\theta) \oplus (\oplus_{v_i \in S_3} K v_i) \oplus (\oplus_{w_j \in S_4} K u_{w_j})$.
- (3) The collection $\{B(S) \mid S \subseteq L_K(E), S \text{ finite}\}$ is an upward directed set of subalgebras of $L_K(E)$.
- (4) $L_K(E) = \varinjlim_{\{S \subseteq L_K(E), S \text{ finite}\}} B(S)$.

Proof (1) By Proposition 1 we have that $F \cup F^* \cup S_1 \cup S_2 \subseteq B(a_1, a_2, \dots, a_\ell)$. Since $S_3 \cup S_4 \subseteq B(a_1, a_2, \dots, a_\ell)$ by construction, (1) follows.

(2) Since the $\{v_i\}$ and $\{u_{w_j}\}$ are pairwise orthogonal idempotents we immediately get that $\sum_{v_i \in S_3} K v_i = \oplus_{v_i \in S_3} K v_i$, and that $\sum_{w_j \in S_4} K u_{w_j} = \oplus_{w_j \in S_4} K u_{w_j}$. We now establish that the three indicated summands are mutually orthogonal, which will establish that the sum $\text{Im}(\theta) + (\oplus_{v_i \in S_3} K v_i) + (\oplus_{w_j \in S_4} K u_{w_j})$ is direct. Let $v \in S_3$. Then by definition v is neither the source vertex nor range vertex for any of the elements in F . In particular, the one dimensional subalgebra Kv of $L_K(E)$ clearly annihilates all the elements of G^0 ; as noted previously, this suffices to yield that Kv indeed annihilates $\text{Im}(\theta)$. But for any $w \in S_4$ we have that $u_w = w - \sum_{f \in F, s(f)=w} f f^*$ is orthogonal in $L_K(E)$ to v , since $S_3 \cap S_4 = \emptyset$.

So we have shown that $\oplus_{v_i \in S_3} K v_i \cap [\text{Im}(\theta) + \oplus_{w_j \in S_4} K u_{w_j}] = \{0\}$. Thus we need only show that $\text{Im}(\theta) \cap K u_w = \{0\}$ for all $w \in S_4$, which we establish by showing

$Ku_w \cdot \text{Im}(\theta) = \text{Im}(\theta) \cdot Ku_w = \{0\}$ and using the fact that each u_w is idempotent. Choose any such w . Since by definition $w \notin r(F)$ we have that $u_w = w - \sum_{f \in F, s(f)=w} ff^*$ is orthogonal in $L_K(E)$ to elements of G^0 of the form $w' - \sum_{g \in F, s(g)=w'} gg^*$ for $w' \in r(F) \cap s(F) \cap s(E^1 \setminus F)$. Similarly, u_w is orthogonal in $L_K(E)$ to $r(e)$ for each $e \in F$. Now suppose $ee^* \in G^0$ with $e \in F$. If $s(e) \neq w$ then $w - \sum_{f \in F, s(f)=w} ff^*$ is clearly orthogonal to ee^* . On the other hand, if $s(e) = w$ then $u_w \cdot ee^* = (w - \sum_{f \in F, s(f)=w} ff^*) \cdot ee^* = wee^* - \sum_{f \in F, s(f)=w} ff^* ee^* = ee^* - ee^* = 0$, with the final simplification occurring because $ff^* ee^* = ee^*$ for $e = f$, and $ff^* ee^* = 0$ otherwise by (CK1). Similarly, we have $ee^* \cdot u_w = 0$. We conclude that the indicated sums are direct.

We now show that the direct sum in fact equals $B(a_1, a_2, \dots, a_\ell)$. By construction, it suffices to show that $u_w \in B(a_1, a_2, \dots, a_\ell)$ for all $w \in S_4$, and that $w \in \text{Im}(\theta) \oplus (\oplus_{v_i \in S_3} Kv_i) \oplus (\oplus_{w_j \in S_4} Ku_{w_j})$ for all $w \in S_4$. But each of these inclusions follow directly by noting that, for each $w \in S_4$ and each $f \in F$ having $s(f) = w$, we have $ff^* \in \text{Im}(\theta)$ by definition.

(3) As noted previously, E_F is a finite graph for each finite subset F of E^1 . In particular, $L_K(E_F)$ is a finitely generated K -algebra (with generating set $E_F^0 \cup E_F^1 \cup (E_F^1)^*$). This in turn implies that $\text{Im}(\theta)$, and hence $B(S)$, is a finitely generated K -algebra for each finite set S of $L_K(E)$. In particular, if S_1 and S_2 are finite subsets of $L_K(E)$, we let T_1 (resp. T_2) denote a finite set of generators of $B(S_1)$ (resp. $B(S_2)$). If $T = T_1 \cup T_2$, then it is clear by construction that $B(S_1) \cup B(S_2) \subseteq B(T)$.

(4) now follows immediately from (1) and (3).

As noted previously, various properties of the graph E need not pass to the graph E_F . However,

Lemma 1 *Let E be any acyclic graph, and F any finite subset of E^1 . Then E_F is acyclic.*

Proof By contradiction, we show that the existence of a closed path in E_F necessarily yields a closed path in E . By definition, a closed path in E_F is of the form $(e_1, e_2), (e_2, e_3), \dots, (e_n, e_1)$ where $(e_i, e_{i+1}) \in E_F^1$. Now it is straightforward to show that the indicated sequence of edges in E_F yields a sequence $e_1, e_2, \dots, e_n$ in E^1 having the desired property.

We recall now some ideas which play central roles in our main result. For additional information about these concepts, see for example [9], [10], and [11].

Definition 4 Let R be a (not necessarily unital) ring.

(1) R is called *von Neumann regular* in case for every $x \in R$ there exists $y \in R$ such that $x = xyx$.

(2) R is called *π -regular* in case for every $x \in R$ there exists $y \in R$ and $n \in \mathbb{N}$ for which $x^n = x^n y x^n$.

(3) R is called *left* (resp. *right*) *π -regular* if for each $a \in R$ there exists $n \in \mathbb{N}$ and $b \in R$ such that $a^n = ba^{n+1}$ (resp. $a^n = a^{n+1}b$). (For rings with local units, this is equivalent to saying that the descending chain of left ideals $Ra \supseteq Ra^2 \supseteq \dots \supseteq Ra^k \supseteq \dots$ (resp. right ideals $aR \supseteq a^2R \supseteq \dots \supseteq a^kR \supseteq \dots$) becomes stationary after finitely many terms.)

(4) R is called *strongly π -regular* if its both left and right π -regular.

Clearly any von Neumann regular ring is π -regular. Conversely, the ring $\mathbb{Z}/4\mathbb{Z}$ provides an easy example of a ring which is π -regular but not von Neumann regular (since $\bar{2}$ has no von Neumann regular inverse).

By [7, Lemma 6], if R is a unital strongly π -regular ring then for every element $a \in R$ there is a positive integer n and an element $x \in R$ such that $ax = xa$ and $a^{n+1}x = a^n = xa^{n+1}$. (We will show below that this result holds for rings with local units as well.) From this property it is then easy to show that if R is strongly π -regular, then R is π -regular. Conversely, the ring $R = \text{End}_K(V)$ of all linear transformations of an infinite dimensional vector space V over a field K provides an example of a ring which is π -regular (in fact, von Neumann regular), but not strongly π -regular. (Indeed, if $\alpha : V \rightarrow V$ is the shift transformation given by $\alpha(K_1) = 0$ and $\alpha(K_{i+1}) = K_i$ for all $i > 1$, then for any n , $\ker \alpha^n = \bigoplus_{i=1}^n K_i$ and so $\alpha^n \neq \beta \alpha^{n+1}$ for any n .)

Lemma 2 *Let R be a ring with local units. Then R is strongly π -regular ring if and only if for every nonzero idempotent v of R , the subring vRv is strongly π -regular.*

Proof Assume R is strongly π -regular. Pick $a \in vRv$. By hypothesis there exists $b \in R$ with $a^n = a^{n+1}b$, and there exists $c \in R$ with $a^m = ca^{m+1}$. But $vav = v$, so $va^n v = a^n$ and $va^{n+1}v = a^{n+1}$. Thus, multiplying both sides of the equation $a^n = a^{n+1}b$ by v , we get $a^n = va^{n+1}v b v = a^{n+1}v b v$. Since $v b v \in vRv$ we have shown that $a^n = a^{n+1}b'$ for some $b' \in vRv$. A similar computation yields that $a^m = c' a^{m+1}$ inside vRv .

Conversely, pick $a \in R$. Then $a \in vRv$ for some idempotent v by definition of set of local units. So there exist b and c in vRv , and hence in R , with the appropriate properties.

Although the properties von Neumann regular, strongly π -regular, and π -regular are in general not equivalent, as one consequence of Theorem 1 we conclude that these properties are indeed equivalent in the context of Leavitt path algebras.

We are now in position to establish our main result.

Theorem 1 *Let E be an arbitrary graph, and let K be any field. The following are equivalent.*

- (1) $L_K(E)$ is von Neumann regular.
- (2) $L_K(E)$ is π -regular.
- (3) E is acyclic.
- (4) $L_K(E)$ is locally K -matricial; that is, $L_K(E)$ is the direct union of subrings, each of which is isomorphic to a finite direct sum of finite matrix rings over K .
- (5) $L_K(E)$ is strongly π -regular.

Proof (1) $\Rightarrow$ (2) is immediate.

(2) $\Rightarrow$ (3). By contradiction, suppose c is a cycle in E , and let $v = s(c) = r(c)$. We show that $v + c$ has no π -regular inverse in $L_K(E)$.

Let γ denote $v + c$. Suppose there exist $\beta \in L_K(E)$, $n \in \mathbb{N}$ such that $\gamma^n \beta \gamma^n = \gamma^n$. Note that, since $\gamma v = \gamma = v \gamma$, we have $\gamma^n v \beta \gamma^n = \gamma^n$. Then $\alpha = v \beta v$ satisfies $\gamma^n \alpha \gamma^n = \gamma^n$ and $v \alpha v = \alpha$. Moreover, $\alpha v = v \alpha = \alpha$.

Write α as a graded sum $\alpha = \sum_{i=M}^N a_i$ where $a_M \neq 0$, $a_N \neq 0$, $\deg(a_i) = i$ for all nonzero a_i having $M \leq i \leq N$, and $a_i = 0$ if $i > N$ or $i < M$. Since $\deg(v) = 0$, the

equation $\alpha v = v\alpha$ implies that $a_i v = v a_i = a_i$ for all i . Now expanding the equation $\gamma^n \alpha \gamma^n = \gamma^n$, we obtain

$$(v + \sum_{k=1}^n \binom{n}{k} c^k) (\sum_{i=M}^N a_i) (v + \sum_{k=1}^n \binom{n}{k} c^k) = v + \sum_{k=1}^n \binom{n}{k} c^k.$$

Equating the lowest degree terms on both sides, we get $va_M v = v$, so that $a_M = v$.

Since $\deg(v) = 0$, we conclude that $M = 0$, and that $a_0 = v$. Thus $\alpha = \sum_{i=0}^N a_i$.

Let $\deg(c) = s > 0$. Now every term other than the first on the right hand side has degree ks for some positive integer $k \leq n$, and so equating the corresponding graded components on both sides, we conclude that $a_i = 0$ if i is not a multiple of s .

We establish by induction that $a_{ks} = f_k(c)$ for each $k \in \mathbb{N}$, where $f_k(c)$ is a polynomial in c with integer coefficients. For $k = 1$, by equating the degree s components on both sides we obtain

$$va_s v + \binom{n}{1} ca_0 + a_0 \binom{n}{1} c = \binom{n}{1} c.$$

This implies that $a_s = -\binom{n}{1}c$, an integral polynomial in c . Now suppose $t > 1$, and suppose $a_{ks} = f_k(c)$, an integral polynomial in c for all $1 \leq k < t$. We expand the previously displayed equation, and equate the degree ts terms of both sides. This yields

$$\begin{aligned} & a_{ts} + \binom{n}{1} c [a_{(t-1)s} + a_{(t-2)s} \binom{n}{1} c + a_{(t-3)s} \binom{n}{2} c^2 + \dots + a_0 \binom{n}{t-1} c^{t-1}] \\ & + \binom{n}{2} c^2 [a_{(t-2)s} + a_{(t-3)s} \binom{n}{1} c + \dots + a_0 \binom{n}{t-2} c^{t-2}] \\ & + \binom{n}{3} c^3 [a_{(t-3)s} + a_{(t-4)s} \binom{n}{1} c + \dots + a_0 \binom{n}{t-3} c^{t-3}] \\ & + \dots + \binom{n}{t} c^t a_0 \\ & = \binom{n}{t} c^t \end{aligned}$$

Substituting for $a_s, \dots, a_{(t-1)s}$ as allowed by the induction hypothesis and solving for a_{ts} , we obtain $a_{ts} = f_t(c)$, a polynomial in c with integer coefficients.

In particular, we conclude that every homogeneous component a_i of α commutes with c in $L_K(E)$. This yields that $c\alpha = \alpha c$. But then the equation $(v+c)^n \alpha (v+c)^n = (v+c)^n$ becomes

$$\alpha(v+c)^{2n} = (v+c)^n.$$

But this is impossible, as follows. Since each a_i is a polynomial in c with integer coefficients, we have $a_i c^r \neq 0$ for all $r \in \mathbb{N}$. Let i be maximal with the property that $a_i(v+c)^{2n} \neq 0$. (Such i exists, since $a_0 = v$ has this property.) Then the left hand side contains terms of degree $2sn + i$ (namely, $a_i c^{2n}$), while the maximum degree of terms on the right hand side is ns .

(3) $\Rightarrow$ (4). We assume E is acyclic. Let $\{B(S) \mid S \subseteq L_K(E), S \text{ finite}\}$ be the collection of subalgebras of $L_K(E)$ indicated in Proposition 2(3). By Proposition 2(4), it suffices to show that each such $B(S)$ is of the indicated form. But by Proposition 2(2), $B(S) = \text{Im}(\theta) \oplus (\oplus_{v_i \in S_3} K v_i) \oplus (\oplus_{w_j \in S_4} K u_{w_j})$. Since terms appearing in the second and third summands are clearly isomorphic as algebras to $K \cong M_1(K)$, it suffices to show that $\text{Im}(\theta)$ is isomorphic to a finite direct sum of finite matrix rings over K . Since E is acyclic, by Lemma 1 we have that E_F is acyclic. But E_F is always finite by definition, so we have by [3, Proposition 3.5] that $L(E_F) \cong \oplus_{i=1}^{\ell} M_{m_i}(K)$ for some $m_1, \dots, m_{\ell}$ in $\mathbb{N}$. Since each $M_{m_i}(K)$ is a simple ring, we have that any homomorphic image of $L_K(E_F)$ must have this same form. So we get that $\text{Im}(\theta) \cong \oplus_{i=1}^L M_{m_i}(K)$ for some $m_1, \dots, m_L$ in $\mathbb{N}$, and we are done. (As remarked previously, since θ is in fact an isomorphism we have $L = \ell$.)

(4) $\Rightarrow$ (1). It is well known that any algebra of the form $\oplus_{i=1}^{\ell} M_{m_i}(K)$ is von Neumann regular. But every element of $L_K(E)$ is contained in a subalgebra of $L_K(E)$ of this form, so that every element of $L_K(E)$ thereby has a von Neumann regular inverse.

(4) $\Rightarrow$ (5). Suppose $L_K(E)$ is locally K -matricial. So every element $a \in L_K(E)$ is contained in a subring $S \cong \oplus_{i=1}^{\ell} M_{m_i}(K)$. As any such S is a unital left (resp. right) artinian ring, there is a $b \in S$ and a positive integer n such $a^n = ba^{n+1}$ (resp. $a^n = a^{n+1}b$).

(5) $\Rightarrow$ (2) By Lemma 2 we have that each $a \in L_K(E)$ is contained in a strongly π -regular unital subring of the form $vL_K(E)v$ for some $v = v^2 \in L_K(E)$. Then by [7, Lemma 6] there is a positive integer n and an element $x \in vL_K(E)v$ such that $ax = xa$ and $a^{n+1}x = a^n = xa^{n+1}$. Now iterating the substitution $a^n = a^{n+1}x = aa^n x = a(a^{n+1}x)x = a^{n+2}x^2$ we get $a^n = a^{2n}x^n$, which using $ax = xa$ gives $a^n = a^n x^n a^n$, which yields (2).

We record the following consequence of Theorem 1, in part because it demonstrates the independence of our results from any cardinality restrictions or graph-theoretic restrictions (e.g. row-finiteness) on the graphs.

Example 3 Let $\aleph$ be any cardinal, and let $\text{Clock}(\aleph)$ be the infinite clock graph having $\aleph$ edges. Then for any field K , the Leavitt path algebra $L_K(\text{Clock}(\aleph))$ is von Neumann regular. In addition, $L_K(\text{Clock}(\aleph))$ is locally K -matricial.

It is worth noting that the locally K -matricial nature of $L_K(\text{Clock}(\aleph))$ does *not* stem from a consideration of the finite complete subgraphs of $\text{Clock}(\aleph)$, since as noted previously $\text{Clock}(\aleph)$ contains *no* such nontrivial subgraphs.

As a second consequence of Theorem 1, we see that the ring $R = \text{End}_K(V)$ of all linear transformations of an infinite dimensional vector space V over a field K cannot be represented as $L_K(E)$ for any graph E , since R is von Neumann regular but not strongly π -regular (as noted earlier). Similarly, let V be a vector space of uncountable dimension over a field K and let S be the (nonunital) subring of $\text{End}_K(V)$ consisting of those linear transformations whose images are of at most countable dimension. Then S is a von Neumann regular ring with local units. However, S is not strongly π -regular, so again invoking Theorem 1 we have that S cannot be represented as the Leavitt path algebra of any graph E .

We conclude this article by analyzing two additional “regularity” properties of a ring. We recall the definitions of some ring-theoretic terms.

Definition 5 Let R be a unital ring.

(1) R is called *clean* if each $a \in R$ is of the form $a = e + u$ where e is an idempotent and u is a (two-sided) unit. If in addition $aR \cap eR = 0$, we say R is a *special clean ring*. A clean ring R is said to be *strongly clean* if in the above definition we can choose e and u which commute.

(2) R is called *unit regular* in case for each $a \in R$ there exists a (two-sided) unit $u \in R$ such that $aua = a$. In particular, every unit regular ring is von Neumann regular.

Additional information about clean rings can be found in [11], while additional information about unit regular rings can be found in [9]. The properties “clean” and “unit regular” are exemplified by matrix rings. Indeed if R is the ring of $n \times n$ matrices over a field, then R is both unit regular [9, page 38] and strongly clean [11, Theorem 4.1]. By [6, Theorem 1], a unital ring R is unit regular if and only if R is a special clean ring; in particular, any ring of the form $M_n(K)$ for K a field and $n \in \mathbb{N}$ is a special clean ring.

While the definitions of von Neumann regularity and π -regularity extend verbatim from unital rings to the nonunital case, the notions of clean and unit regularity require additional attention in the nonunital situation (since each definition refers to a unit in the given ring). We now show how to naturally extend these latter two notions to rings with local units.

Definition 6 Let R be a ring with local units.

(1) R is called *locally unit regular* if for each $a \in R$ there is an idempotent $v \in R$ for which $a \in vRv$, and elements $u, u' \in vRv$ such that $uu' = v = u'u$, and $aua = a$.

(2) R is called *locally clean* if for each $a \in R$ there is an idempotent $v \in R$ for which $a \in vRv$, and elements $e, u, u' \in vRv$ such that e is an idempotent, $uu' = v = u'u$, and $a = e + u$.

That the two notions given in the previous definition are natural generalizations of the corresponding notions for unital rings is established in the following.

Lemma 3 Let R be a unital ring.

- (1) R is locally unit regular if and only if R is unit regular.
- (2) R is locally clean if and only if R is clean.

Proof For (1), suppose R is a ring with 1 and is locally unit regular. Let $a \in R$, and let v, u, u' as given in the definition. Then $w = u + (1 - v)$ and $w' = u' + (1 - v)$ satisfy $ww' = 1 = w'w$ and $a = awa$. Hence R is unit regular. The converse is clear with $v = 1$.

Likewise, for (2), suppose R is a ring with 1 and is locally clean. Let $a \in R$, and write $a = u + e$ as given in the definition. Then $a = w + e'$, where $e' = e + (1 - v)$ is an idempotent and $w = u - (1 - v)$ is a two-sided unit in R (since with $w' = u' - (1 - v)$, we have $ww' = 1 = w'w$). Thus R is clean. As with (1), the converse follows with $v = 1$.

Our final result shows that for acyclic graphs E , $L_K(E)$ possesses the locally unit regular property, as well as a property involving clean unital subrings.

Theorem 2 *Let E be an arbitrary graph, and let K be any field. Then the following conditions are equivalent:*

- (1) *E is acyclic.*
- (2) *$L_K(E)$ is locally unit regular.*
- (3) *$L_K(E)$ is a direct limit of unital strongly clean rings, each of which is special.*

Proof (1) $\Leftrightarrow$ (2) Suppose E is acyclic. Then, by Theorem 1, $L_K(E)$ is a direct union of direct sums of matrix rings each of which, by [9, page 38], is unit regular. It is then clear that $L_K(E)$ is locally unit regular, where for each $a \in L_K(E)$ we use for v the identity element of the corresponding subring $B(S)$. Conversely, if $L_K(E)$ is locally unit regular, then it is, in particular, von Neumann regular. So, by Theorem 1, E is acyclic.

(2) $\Leftrightarrow$ (3) Suppose $L_K(E)$ is locally unit regular. Since it is von Neumann regular, Theorem 1 implies that $L_K(E)$ is a directed union of direct sums of matrix rings L_i each of which, as noted above, is a special clean ring which is, in addition, strongly clean. On the other hand, if $L_K(E)$ is a directed union of special clean rings L_i , then each L_i is unit regular by [6, Theorem 1], and so $L_K(E)$ is locally unit regular. (Again for each $a \in L_K(E)$ we use for v the identity element of the corresponding subring $B(S)$.)

A study of $L_K(E)$ for arbitrary graphs E is presented by Goodearl in [8]. Included in [8] is a method to write E as a direct union of countable complete subgraphs. We now show how this approach together with the desingularization process yields an alternate proof of the implication (3) $\Rightarrow$ (1) of Theorem 1. Our aim in doing so is to contrast this approach with that of using Proposition 1, which helps us to establish not only the von Neumann regularity of $L_K(E)$ for an acyclic graph E , but uncovers several internal properties of such an $L_K(E)$ (e.g., locally matricial and locally unit regular). (Our approach also shows the coincidence of von Neumann regularity with π -regularity and strong π -regularity for Leavitt path algebras.) One may also note that the desingularization approach as shown below does not work for π -regular rings since π -regularity, unlike von Neumann regularity, is not a Morita invariant (see e.g. [12]). Proposition 1 poses no such restrictions, and provides additional structural insight into these rings.

So suppose E is acyclic. By [8, Proposition 2.7] $L_K(E) = \varinjlim_{\alpha \in A} L_K(E_\alpha)$, with the limit taken over the set $\{E_\alpha \mid \alpha \in A\}$ of countable complete subgraphs of E . So in order to show that $L_K(E)$ is von Neumann regular, it suffices to show that each $L_K(E_\alpha)$ is von Neumann regular, since the direct limit of von Neumann regular rings is von Neumann regular. Since E is acyclic then necessarily so is each E_α .

Since E_α is countable, we may form a desingularization F_α of E_α . (See e.g. [2].) By construction, F_α is row-finite. Also, since desingularization preserves Morita equivalence, and von Neumann regularity is preserved by Morita equivalence for rings with local units by [4, Proposition 3.1], it suffices to show that each $L_K(F_\alpha)$ is von Neumann regular. Since each E_α is acyclic, the desingularization construction shows that each F_α is acyclic as well.

But by [5, Lemma 3.2] F_α is the direct union of G_β (the union taken over the set of finite complete subgraphs of F_α), and $L_K(F_\alpha) = \varinjlim_{\beta \in B} L_K(G_\beta)$. Thus it suffices to show that each $L_K(G_\beta)$ is von Neumann regular. Since G_β is a subgraph of F_α we have that G_β is acyclic.

So in the end, to establish that $L_K(E)$ is von Neumann regular, it suffices to show that for any finite acyclic graph G that $L_K(G)$ is von Neumann regular. But by [3, Proposition 3.5] the Leavitt path algebra of a finite acyclic graph is isomorphic to a finite direct sum of finite dimensional matrix rings over the ground field K , and such rings are well known to be von Neumann regular (see e.g. [9, Section 1]).

We conclude this article by noting one more consequence of Theorem 1 (we thank the referee for this suggestion). The proof follows directly from the fact that von Neumann regularity is a Morita invariant for rings with local units. We contrast this result with the aforementioned remark that, in general, the π -regularity property is not a Morita invariant.

Corollary 1 *The property of π -regularity is a Morita invariant for Leavitt path algebras; that is, if E and F are graphs with $L_K(E)$ Morita equivalent to $L_K(F)$, then $L_K(E)$ is π -regular if and only if $L_K(F)$ is π -regular, and in this case E and F are both acyclic.*

acknowledgments

The authors thank E. Pardo and M. Siles Molina for their valuable discussions during the preparation of this paper. The authors also thank the referee for a very careful reading of, and suggested changes to, the initial version of the manuscript.

References

1. ABRAMS, G., ÁNH, P.N., LOULY, A., PARDO, E.: The classification question for Leavitt path algebras, *J. Algebra*, to appear. ArXiv: 0706.3874.
2. ABRAMS, G., ARANDA PINO, G.: The Leavitt path algebra of arbitrary graphs, *Houston J. Math.*, to appear.
3. ABRAMS, G., ARANDA PINO, G., SILES MOLINA, M.: Finite-dimensional Leavitt path algebras, *J. Pure Appl. Algebra*. **209**(3), 753–762 (2007).
4. ÁNH, P.N., MÁRKI, L.: Morita equivalence for rings without identity, *Tsukuba J. Math.* **11**(1), 1–16 (1987).
5. ARA, P., MORENO, M.A., PARDO, E.: Nonstable K-Theory for graph algebras, *Algebra Represent. Theory* **10**(2), 157–178 (2007).
6. CAMILLO, V., KHURANA, D.: A characterization of unit regular rings, *Comm. Algebra* **29**(5), 2293–2295 (2001).
7. CAMILLO, V., YU, H.-P.: Stable range one for rings with many idempotents, *Trans. A.M.S.* **347**(8), 3141–3147 (1995).
8. GOODEARL, K.: Leavitt path algebras and direct limits, to appear. arXiv:0712.2554v1
9. GOODEARL, K.: *Von Neumann Regular Rings*. Krieger Publ., Malabar, FL (1991). ISBN 0-89464-632-X.
10. KAPLANSKY, I.: Topological representation of algebras. II, *Trans. A.M.S.* **68**(1), 62–75 (1950).
11. NICHOLSON, W.K., ZHOU, Y.: Clean Rings: a survey. In: *Advances in Ring Theory: Proceedings of the 4th China-Japan-Korea International Conference*, pp. 181–198. World Sci. Publ., Hackensack, N.J. (2005). ISBN: 981-256-425-X.
12. ROWEN, L.: Examples of semiperfect rings, *Israel J. Math.* **65**(3), 273–283 (1989).
13. RAEBURN, I., SZYMAŃSKI, W.: Cuntz-Krieger algebras of infinite graphs and matrices, *Trans. A.M.S.* **356**(1), 39–59 (2003).